

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)  
(USPTO) THIS PAGE BLANK (USPTO)



**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re: application of  
**Mercken et al.**

Examiner: **To Be Assigned**

Group Art Unit: **1614**

Application No.: **10/691,079**

Filed: **October 22, 2003**

Title: **INHIBITORS OF SRC KINASE FOR USE IN  
ALZHEIMER'S DISEASE**

**CERTIFICATE OF MAILING (37 CFR 1.8a)**

I hereby certify that this paper (along with any referred to as being attached or enclosed) is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner for Patents, P. O. Box 1450, Alexandria, VA 22313-1450.

Date of Deposit

May 10, 2004

Delia Coughlin

(Type or print name of person mailing paper)

Delia Coughlin

(Signature of person mailing paper)

Mail Stop  
Commissioner for Patents  
P. O. Box 1450  
Alexandria, VA 22313-1450

**SUBMISSION AND REQUEST FOR ENTRY  
OF PRIORITY PAPERS 37 C.F.R. § 1.55(a)**

Applicants submit herewith a certified copy of EP application 02292608.3, filed on October 22, 2002, for which priority is claimed in the above-identified application.

This submission and request for entry is being made to satisfy the requirements under 35 U.S.C. § 119. Please note that no fees are associated with the entry of the priority documents since they are being timely submitted prior to the date the issue fee is due.

Respectfully submitted,

Karen I. Krupen

Karen I. Krupen, Reg. No. 34,647  
Attorney/Agent for Applicant

Aventis Pharmaceuticals Inc.  
Patent Department  
Route #202-206 / P.O. Box 6800  
Bridgewater, New Jersey 08807-0800  
Telephone (908) 231-4658  
Telefax (908) 231-2626

Aventis Docket No. **FRAV2002/0030USNP**





**Europäisches  
Patentamt**

**European  
Patent Office**

**Office européen  
des brevets**

**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

**Patentanmeldung Nr.    Patent application No.    Demande de brevet n°**

02292608.3

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**R C van Dijk**

THIS PAGE BLANK (USPTO)  
THIS PAGE BLANK (USPTO)



Anmeldung Nr:  
Application no.: 02292608.3  
Demande no:

Anmeldetag:  
Date of filing: 22.10.02  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Aventis Pharma S.A.  
20, avenue Raymond Aron  
92160 Antony  
FRANCE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Inhibitors of Src kinase for use in Alzheimer's disease

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

G01N33/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

THIS PAGE BLANK (USPTO)



### Inhibitors of Src kinase for use in Alzheimer's disease

The present invention relates to the identification of inhibitors of Src kinase, assays and methods useful for such identification and the use of Src inhibitors for the preparation of pharmaceuticals for the treatment of Alzheimer's disease.

Alzheimer's disease is a neurodegenerative disorder of the brain, which is accompanied at the cellular level by a massive loss of neurons in the limbic system and in the cerebral cortex. In the brain areas affected, protein deposits, so-called plaques, can be detected at the molecular level, which are an essential characteristic of Alzheimer's disease. The protein occurring most frequently in these plaques is a peptide of 40 to 42 amino acids in size, which is designated as A $\beta$ -peptide. This peptide is a cleavage product of a larger protein of 695 to 770 amino acids, the so-called Amyloid Precursor Protein (APP).

APP is an integral transmembrane protein, which firstly traverses the lipid bilayer. By far the largest part of the protein is extracellular, while the shorter C-terminal domain is directed into the cytosol (Figure 1). APP is the substrate of three different proteases:  $\alpha$ -secretase,  $\beta$ -secretase and  $\gamma$ -secretase. Within APP, about two thirds of the A $\beta$ -peptide originates from the extracellular domain and about one third from the transmembrane domain.

Beside the membrane-linked APP, a secreted form of APP can be detected which consists of the large ectodomain of APP and is designated as APP<sub>sec</sub> ("secreted APP"). APP<sub>sec</sub> is formed from APP by proteolytic cleavage, which is effected by the  $\alpha$ -secretase (Figure 2). This proteolytic cleavage takes place within the amino acid sequence of the A $\beta$ -peptide (after amino acid residue 16 of the A $\beta$ -peptide). Proteolysis of APP by the  $\alpha$ -secretase thus excludes the formation of the A $\beta$ -peptide.

The A $\beta$ -peptide can thus only be formed from APP in an alternative processing route. It is postulated that two proteases are involved in this processing, one protease, which is designated as  $\beta$ -secretase, cleaving at the N-terminus of the A $\beta$ -peptide in the APP and the second protease, which is designated as  $\gamma$ -secretase, releasing the C-terminus of the A $\beta$ -peptide (Kang, J. et al., (1987) Nature, 325, 733 (Figure 2).

There are many indications that the A $\beta$ -peptide is a crucial factor in the occurrence of Alzheimer's disease. Inter alia, neurotoxicity of A $\beta$ -fibrils in cell culture is postulated (Yankner, B.A. et al., (1990) Proc Natl Acad Sci USA, 87, 9020). In patients with Down's syndrome, in which APP occurs in an additional copy, the neuropathology characteristic of Alzheimer's disease also occurs even at an age of 30 years. Here, it is assumed that the overexpression of APP follows an increased conversion into the A $\beta$ -peptide (Rumble, B. et al., (1989), N. Engl. J. Med., 320, 1446).

Furthermore the strongest indication of the central role of the A $\beta$ -peptide in Alzheimer's disease are the familial forms of the disease. Here, mutations are found in the APP gene around the area of the secretase cleavage sites or in two further AD-associated genes (presenilins), which in cell culture lead to a significant increase in A $\beta$  production (Scheuner, D. et al., (1996), Nature Medicine, 2, 864).

There are a number of indications that APP is first cleaved by the  $\beta$ -secretase, to serve thereafter as a substrate for  $\gamma$ -secretase (Maruyama, K. Y. et al., (1994) Biochem. Biophys Res Commun, 202, 1517; Estus, S. et al., (1992), Science, 255, 726 ). The  $\gamma$ -secretase therefore has a crucial role in the formation of the A $\beta$ -peptide. A demonstration of the activity of the  $\gamma$ -secretase which is customarily used is the detection of the A $\beta$ -peptide, which, however, frequently turns out to be difficult.

An important reason for this is that only a small part of APP is converted into the A $\beta$ -peptide (Simons M, et al., *Neurosci* (1996) 1;16(3):899-908). Moreover, the A $\beta$ -peptide is a small breakage fragment of about 4 kDa and,  
 5 on account of its hydrophobic character, has a great tendency to self-aggregation so that it easily precipitates under physiological conditions (Hilbich, C. et al., (1991) *J. Mol. Biol.*, 218, 149).

The short, 47 amino acid-long C-terminal tail of APP is exposed to the cytosol  
 10 and is the site of interaction for molecular adaptors containing PTB domains, named Fe65, X11, m-dab and Jip1. Since the binding of these proteins is dependent on the YENPTY sequence on APP, it is expected that their interactions are not simultaneous (T. Russo et al. (1998) *FEBS Lett.* 434, 1-7; N. Zambrano et al. (1997) *J. Biol. Chem.* 272, 6399-6405). Indeed, it has  
 15 been shown, in cultured cells, that Fe65 and X11 proteins have opposite effects on the proteolytic processing of APP (Sabo S. et al. (1999) *J. Biol. Chem.* 274, 7952-7957; Borg J.P. et al. (1998) *J. Biol. Chem.* 273, 14761-14766). A possible explanation for these findings may reside in the recruitment, by either Fe65 or X11, of APP in different macromolecular  
 20 complexes, depending on the interaction of different sets of proteins with the other protein-protein interaction domains of the two adaptors.

The APP cytodomain contains various serine and threonine residues, which are phosphorylated *in vitro* by different kinase activities. Thr668 is a main site  
 25 of phosphorylation *in vivo*, being phosphorylated by neuronal cyclin-dependent kinases, Cdk5 in neurons (Iijima, K. et al. (2000) *J. Neurochem.* 75, 1085-1091), cdc2 kinase in dividing cells (Suzuki, T. et al. (1994) *EMBO J.* 13, 1114-1122; Oishi, M. et al. (1997) *Mol. Med.* 3, 111-123), and glycogen synthase kinase 3b (GSK3b) and stress-activated protein kinase 1b  
 30 (SAP kinase1b) *in vitro* (Aplin, E. E. et al. (1996) *J. Neurochem.* 67, 699-707; Standen, C. L. et al. (2001) *J. Neurochem.* 76, 316-320). It has been shown

that phosphorylation of APP regulates neurite extension in differentiating PC12 cells (Ando, K. et al. (1999) *J. Neurosci.* 19, 4421-4427), while, at the molecular level, APP phosphorylation may regulate the binding to the PTB domain-containing adaptors and its processing (Ando K. et al. (2001) *J Biol. Chem.*, 276, 40353-40361).

Previously it was demonstrated that APP is phosphorylated at tyrosine 682 in cells expressing an active form of the Abl non receptor tyrosine kinase (Zambrano N. et al. (2001) *J Biol. Chem.* 276, 19787-92); active Abl is recruited in proximity to APP by Fe65, which may bind simultaneously these two proteins through its WW domain (Abl) and PTB2 domain (APP). Phosphorylation of Tyr682 of APP generates a docking site for the SH2 domain of Abl and, in fact stable complexes between APP and Abl are formed.

In order to test the hypothesis that tyrosine phosphorylation regulates pp60c-src biological activity, Kmiecik and Shalloway (1987) *Cell* 49, 65-73 described the construction and study of pp60c-src mutants in which Tyr 527 and Tyr 416 were separately or coordinately altered to phenylalanine. Tyr----Phe 527 mutation strongly activated pp60c-src transforming and kinase activities, whereas the additional introduction of a Tyr----Phe 416 mutation suppressed these activities. Tyr----Phe 416 mutation of normal pp60c-src eliminated its partial transforming activity, which suggests that transient or otherwise restricted phosphorylation of Tyr 416 is important for pp60c-src function even though stable phosphorylation is not observed in vivo. Normally, pp60c-src is phosphorylated in vivo at tyrosine 527, a residue not present in pp60v-src (its transforming homolog), and not at tyrosine 416, its site of in vitro autophosphorylation.

Williamson et al. (2002) *J. Neurochem.* 22, 10-20) described the rapid phosphorylation of neuronal proteins including Tau and Focal adhesion

kinase in response to amyloid- $\beta$  peptide exposure and an involvement of Src family protein kinases.

Slack, B. E. and Berse; B. (1998) described in Society for Neuroscience Abstracts, Vol. 24, No. 1-2, pp. 208 (Conference/Meeting Information: 28<sup>th</sup> Annual Meeting of the Society for Neuroscience, Part 1, Los Angeles, California, USA, Nov. 7-12, 1998 Society for Neuroscience, ISSN: 0190-5295) a role for tyrosine kinases in the stimulation of APP release by action of muscarinic m3 acetylcholine receptors. They described a role for Src tyrosine kinase in the regulation of APP<sub>sec</sub> release by muscarinic receptors. They demonstrated that an increase in the active form of Src leads to a decrease in secAPP.

The present invention provides methods of identifying therapeutic agents for the treatment of Alzheimer's disease.

One embodiment of the invention provides methods for identifying inhibitors of Src activity, whereby the methods comprise the steps

- a) providing a Src protein (i.e. a Src encoding DNA under control of an expression element) and
- b) determining if a compound inhibits the activity of Src.

The Src protein could be any mammalian Src protein, preferably human Src (SEQ ID NO. 1 (isoform 1) and SEQ ID NO. 2 (isoform 2)) or rodent Src (SEQ ID NO. 3). The protein could be obtained by expressing human Src or rodent Src cDNA, by using e. g. sequences SEQ ID NO. 4 or SEQ ID NO. 5. In addition, one of the sequences deposited under Genbank Accession No. M17031 or BC011566 can be used.

For the method preferably a mammalian cell or cell line is used in which Src is expressed, either naturally or because the cell / cell line is genetically engineered. In a particular embodiment primary cultures of neurones are used.

5

Another embodiment of the invention relates to a method of identifying compounds, which inhibit Src expression. The method comprises the steps

- a) providing a sequence which regulates Src expression (i.e. a Src  
10           promotor sequence) and
- b) determining if a compound inhibits the expression of Src protein.

The sequence which regulates Src expression is preferably linked to a reporter gene or a reporter gene construct. Such reporter gene can easily be  
15       used to determine, if a compound inhibits the expression of Src. Another possibility is the use of the Src gene. The region of chromosome 20 including the Src gene is deposited under Genbank Accession no AL133293 .

An house-keeping promoter, which can be used as a sequence which  
20       regulates Src expression, has been described, as well as an alternative promoter regulated by the Hepatic Nuclear Factor-1a (Bonham et al. (2000) J. Biol. Chem. 275, 37604) (Genbank accession number AF272982).

Active compounds, this means compounds, which inhibit Src expression or  
25       which inhibit Src protein activity, can be used as pharmaceuticals. Such compounds can be used for the preparation of a pharmaceutical for the treatment or prevention of Alzheimer's disease.

Another embodiment of the invention relates to the use of compounds identified by one of the forgoing methods for preparing a pharmaceutical for  
30       the treatment of Alzheimer's disease.

The invention further relates to a method of preparing a pharmaceutical for treatment of Alzheimer's disease comprising the steps

- 5 a) identification of a therapeutic compound by the use of one of the methods described above;
- b) optimisation of the identified compound; and
- c) formulation of the optimised compound.

10 Another embodiment of the invention relates to the use of Src inhibitors known in the art for the preparation of a pharmaceutical for the treatment or the prevention of Alzheimer's disease. The invention relates to the use of PP2 (4-amino-5-(4-chlorophenyl)-7-(t-butyl)pyrazolo[3,4-d]pyrimidine) as pharmaceutical. The invention further relates to the use of PP2 for the preparation of a pharmaceutical for the treatment of Alzheimer's disease.

15

Another embodiment of the invention refers to the use of PP2 for the preparation of a pharmaceutical for the treatment of Alzheimer's disease.

20 The new results demonstrate, that APP is phosphorylated not only by Abl. In fact Src kinase is also responsible for APP tyrosine phosphorylation. It is expected that tyrosine phosphorylation of APP alters the binding properties of its cytodomain, and then the processing of APP could be consequently modified. If so, this will prove useful the use of currently available tyrosine kinase inhibitors, and to design new ones for pharmacological modulation of

25 the release of APP processing products in cellular and animal models of Alzheimer's disease.

Taken together, the results from the following examples suggest that exogenous Src activity is responsible for activation of a pathway resulting in increased A $\beta$  secretion. This also suggests that PP2 could be used as a

30 pharmacological agent able to reduce A $\beta$  secretion in cellular and animal systems resembling AD status.

Therefore, a line of CHO cells stably overexpressing APP751 form was used, which secretes discrete levels of A $\beta$ . If secretion involves endogenous Src activity, PP2 should be able to inhibit production of A $\beta$  also in this cellular system. This is indeed the case since, as shown in Figure 6, PP2 reduces in a dose-dependent manner the amount of A $\beta$  at the three time points tested (1, 3 and 12 hours), while PP3, used at the highest concentration, has no effect.

These experiments clearly outline the relevance of Src activation in A $\beta$  secretion and constitute the basis to try pharmacological applications for tyrosine kinase inhibitors in the treatment of AD. Screening of such inhibitors and their optimization will lead to the development of potential, innovative therapeutic agents for treatment of Alzheimer's disease.

## 15 Examples

### Example 1

#### Cell culture, transfections and treatments

20 Human embryonic kidney cells (HEK293) were cultured in Dulbecco's modified minimal medium supplemented with 10% fetal calf serum at 37 °C in a 5% CO<sub>2</sub> atmosphere. For transfection, cells were grown for 16 hours in antibiotic-free medium, and transfected with Lipofectamine 2000 (Invitrogen) according to manufacturer's instructions. The total amount of DNA was maintained constant by addition of empty vector DNA. 48 hours after transfection, culture medium were recovered, while cells were harvested in ice-cold phosphate buffer saline (PBS) and gently lysed in lysis buffer (50 mM Tris/HCl, pH 7.5, 150 mM NaCl, 0.5% Triton X-100, 10% glycerol, 50 mM NaF, 1mM Na vanadate), in the presence of a protease inhibitor cocktail (Complete, EDTA-free, Roche). The extracts were clarified by centrifugation at 16,000 x g at 4 °C, and the protein concentration determined by the Bio-



Rad protein assay according to manufacturer's instructions. For metabolic labelling, 36 hours after transfection, cells were incubated for 30 minutes in medium without methionine and cysteine, then the medium was replaced with fresh medium containing 80  $\mu\text{Ci/ml}$  of  $^{35}\text{S}$ -methionine/cysteine mixture (Promix, 1000 Ci/mmol, Amersham Pharmacia Biotech). Pulse was for 30 minutes, then the radioactive medium was replaced with complete medium for 90 minutes. CHO cells expressing wild-type APP751 have been described (N. Zambrano et al. (1997) J. Biol. Chem. 272, 6399-6405).

Immuno-precipitations were performed either on culture medium, or on cellular lysates with 1  $\mu\text{g/sample}$  of the 6E10 monoclonal antibody (Signet). Radioactive samples were resolved on 10% SDS-PAGE and quantitated with a Storm 840 phosphorimager (Amersham Pharmacia Biotech).

PP2 and PP3 (Hanke, J.H. et al. (1996) J. Biol. Chem. 271, 695–701) were purchased from Calbiochem and dissolved in DMSO; treatments were performed on transfected cells for 48 hours with vehicle alone, or with 1, 5 and 20  $\mu\text{M}$  PP2 or PP3. Concentration of DMSO was the same in all samples. Treatments on CHO/APP751 cells was made with DMSO, or PP2 (5 and 20  $\mu\text{M}$ ), or PP3 (20  $\mu\text{M}$ ).

## Example 2

HEK293 cells were transfected with APP695 expression vectors. This cellular system was used because these cells are transfected to high efficiency, and because they express the whole set of processing activities required for APP maturation. In fact, upon transfection of human APP695 expression vectors, both  $\alpha$ -secretase product ( $\text{APP}_{\text{sec}}$ ) and  $\beta$ -  $\gamma$ -secretase products ( $\text{A}\beta$ ) accumulate in the culture medium. HEK293 cells were co-transfected with APP and either empty vector, or active Abl (Abl-PP), or active Src (SrcYF) kinases, and the secretion of  $\text{APP}_{\text{sec}}$  and  $\text{A}\beta$  were evaluated. Transfected cells have been pulse-labelled with  $^{35}\text{S}$ -Met/Cys mix for 30', and chased with cold amino acids for 90'. Both medium and protein extracts have been used

in immunoprecipitation experiments for quantitative dosage of radiolabeled APP<sub>sec</sub> and holo-APP, respectively. While the stability of holo-APP was not affected by co-transfection with either Abl-PP or SrcYF active kinases (Figure 3A), a reduction in the relative amounts of APP<sub>sec</sub>, normalised to the  
 5 corresponding holo-APP synthesised during the pulse period was observed in both cases (Figure 3B). This can be interpreted as a decrease of the activity of the  $\alpha$ -secretase pathway upon co-transfection of APP with the active kinases.

### 10 Example 3

Next, the accumulation of A $\beta$  from cells transfected as above was investigated. The dosage was performed by sandwich ELISA assays on culture medium at 48 hours after transfection (Figure 4). The assay clearly  
 15 shows that in Src-expressing cells the amount of A $\beta$  secreted is more elevated than in the presence of the control vector, or Abl-PP.

The mechanism by which active Src elicits such a dramatic increase in A $\beta$  secretion is not clear. However, it might depend on phosphorylation of some other proteins, different from APP, since tyrosine phosphorylation of APP by  
 20 Abl-PP does not result in increased A $\beta$  levels. Accordingly, a similar behaviour is obtained with various APP mutants bearing tyrosine substitutions to either glycine or phenylalanine (data not shown).

Anyway, the rise in A $\beta$  levels does depend on the tyrosine kinase activity of Src, since the accumulation of A $\beta$  is sensitive to treatment by the Src-family specific inhibitor PP2. In fact, as shown in Figure 5, the exposure of  
 25 transfected cells for 48 hours to increasing concentrations of PP2 results in dose-dependent decrease of secreted A $\beta$ . Either PP3, an inactive analog of PP2, or vehicle alone, do not affect A $\beta$  rise by SrcYF transfection.

Taken together, these observations suggest that exogenous Src activity is responsible for activation of a so far unidentified pathway resulting in increased A $\beta$  secretion.

## 5 REFERENCES

- 1) T. Russo et al. (1998) FEBS Lett. 434, 1-7.
- 2) N. Zambrano et al. (1997) J. Biol. Chem. 272, 6399-6405.
- 3) Sabo S. et al. (1999) J. Biol. Chem. 274, 7952-7957.
- 4) Borg J.P. et al. (1998) J. Biol. Chem. 273, 14761-14766.
- 10 5) Iijima, K. et al. (2000) J. Neurochem. 75, 1085-1091.
- 6) Suzuki, T. et al. (1994) EMBO J. 13, 1114-1122.
- 7) Oishi, M. et al. (1997) Mol. Med. 3, 111-123.
- 8) Aplin, E. E. et al. (1996) J. Neurochem. 67, 699-707.
- 9) Standen, C. L. et al. (2001) J. Neurochem. 76, 316-320.
- 15 10) Ando, K. et al. (1999) J. Neurosci. 19, 4421-4427.
- 11) Ando K. et al. (2001) J Biol Chem.;276, 40353-40361.
- 12) Zambrano N. et al. (2001) J Biol Chem 276, 19787-19792.
- 13) Hanke, J.H. et al. (1996) J. Biol. Chem. 271, 695-701.

20

Table 1:

Protein sequence of isoform 1 of human Src protein (SEQ ID NO. 1)

MGSNKS PKDASQRRRSLEPAENVHGAGGGAFPASQTPSKPASADGHRGPSAAFAPAAA  
 25 EPKLFGGFNSSDVTVTSPQRAGPLAGGVTTFVALYDYESRTETDLSFKKGERLQIVNNTRKVD  
 VREGDWWLAHSLSTGQTYIPSNYVAPSDSIQAEWYFGKITRRESERLLLNAENPRGTFL  
 VRESETTKGAYCLSVSDFDNAKGLNVKHYKIRKLDSGGFYITSRTQFNSLQQLVAYYSKHAD  
 GLCHRLTTVCPTSKPQTQGLAKDAWEIPRESLRLEVKLGGQCFGEVWMGTWNGTTRVAIK  
 TLKPGTMSPEAFLQEAQVMKKLRHEKLVQLYAVVSEEPYIVTEYMSKGSLLDFLKGETGKYL

RLPQLVDMAAQIASGMAYVERMNYVHRDLRAANILVGENLVCKVADFGLARLIEDNEYTARQ  
 GAKFPIKWTAPEAALYGRFTIKSDVWSFGILLTELTTKGRVPYPGMVNREVLDQVERGYRMP  
 CPPECPESLHDLMCQCWRKEPEERPTFEYLQAFLEDYFTSTEPQYQPGENL

5 Table 2:

Protein sequence of isoform 2 of human Src (SEQ ID NO. 2)

MGSNKS PKDASQRRRSLEPAENVHGAGGGAFFASQTPSKPASADGHRGPSAAFAPAAA  
 EPKLFGGFNSSDTVTSPQRAGPLAGGVTTFVALYDYESRTETDLSFKKGERLQIVNNTGED  
 10 WWLAHSLSTGQGTGYIPSNYVAPSDSIQAEWYFGKITRRESERLLLNAENPRGTFLVRESET  
 TKGAYCLSVSDFDNAKGLNVKHYKIRKLDSSGGFYITSRTQFNSLQQLVAYYSKHADGLCHRL  
 TTVCPSTSKPQTQGLAKDAWEIPRESLRLEVKLGGQCFGEVWMGTWNGTTRVAIKTLKPGT  
 MSPEAFLQEAQVMKKLRHEKLVQLYAVVSEPIYIVTEYMSKGSLLDFLKGETGKYLRPLPQL  
 VDMAAQIASGMAYVERMNYVHRDLRAANILVGENLVCKVADFGLARLIEDNEYTARQGAKF  
 15 PIKWTAPEAALYGRFTIKSDVWSFGILLTELTTKGRVPYPGMVNREVLDQVERGYRMPCPPE  
 CPESLHDLMCQCWRKEPEERPTFEYLQAFLEDYFTSTEPQYQPGENL

Table 3:

Sequence of mouse Src protein (SEQ ID NO. 3)

20

MGSNKS PKDASQRRRSLEPSENVHGAGGGAFFASQTPSKPASADGHRGPSAAFVPPAAEP  
 KLFGGFNSSDTVTSPQRAGALAGGVTTFVALYDYESRTETDLSFKKGERLQIVNNTRKVDVR  
 EGDWWLAHSLSTGQGTGYIPSNYVAPSDSIQAEWYFGKITRRESERLLLNAENPRGTFLVR  
 ESETTKGAYCLSVSDFDNAKGLNVKHYKIRKLDSSGGFYITSRTQFNSLQQLVAYYSKHADGL  
 25 CHRLTTVCPTSKPQTQGLAKDAWEIPRESLRLEVKLGGQCFGEVWMGTWNGTTRVAIKTLK  
 PGTMSPEAFLQEAQVMKKLRHEKLVQLYAVVSEPIYIVTEYMNKGSLLDFLKGETGKYLRPL

PQLVDMSAQIASGMAYVERMNYVHRDLRAANILVGENLVCKVADFGLARLIEDNEYTARQG  
 AKFPIKWTAPEAALYGRFTIKSDVWSFGILLTELTTKGRVPYPGMVNREVLQDQVERGYRMPC  
 PPECPESLHDLMCQCWRKEPEERPTFEYLQAFLEDYFTSTEPQYQPGENL

5 Table 4:

cDNA sequence encoding human Scr protein (SEQ ID NO. 4)

1 catcgagggtt ttgagaggct aactctccca aaaaggacca tgggtagcaa  
 10 caagagcaag  
 61 cccaaggatg ccagccagcg gcgcgcgagc ctggagcccg ccgagaacgt  
 gcacggcgct  
 121 ggcggggggcg ctttccccgc ctgcagacc ccagcaagc cagcctcggc  
 cgacggccac  
 15 181 cgcgggccca gcgcggcctt cgcccccgcg gccgcgagc ccaagctgtt  
 cggaggcttc  
 241 aactcctcgg acaccgtcac ctccccgcag agggcgggccc cgctggccgg  
 tggagtgacc  
 301 acctttgtgg ccctctatga ctatgagtct aggacggaga cagacctgtc  
 20 cttcaagaaa  
 361 ggcgagcggc tccagattgt caacaacaca gagggagact ggtggctggc  
 ccactcgctc  
 421 agcacaggac agacaggcta catccccagc aactacgtgg cgccctccga  
 ctccatccag  
 25 481 gctgaggagt ggtatttttg caagatcacc agacgggagt cagagcggtt  
 actgctcaat  
 541 gcagagaacc cgagagggac cttcctcgtg cgagaaagtg agaccacgaa  
 aggtgcctac  
 601 tgcctctcag tgtctgactt cgacaacgcc aagggcctca acgtgaagca  
 30 ctacaagatc  
 661 cgcaagctgg acagcggcgg cttctacatc acctcccgca ccagttcaa  
 cagcctgcag  
 721 cagctgggtg cctactactc caaacacgcc gatggcctgt gccaccgcct  
 caccaccgtg  
 35 781 tgccccacgt ccaagccgca gactcagggc ctggccaagg atgcctggga  
 gatccctcgg

841. gagtcgctgc ggctggaggt. caagctgggc. cagggtctgt ttggcgaggt.  
 gtggatgggg  
 901 acctggaacg gtaccaccag ggtggccatc aaaacctga agcctggcac  
 gatgtctcca  
 5 961 gaggccttcc tgcaggaggc ccaggtcatg aagaagctga ggcattgagaa  
 gctgggtgcag  
 1021 ttgtatgctg tggtttcaga ggagccatt tacatcgtca cggagtacat  
 gagcaagggg  
 1081 agtttgctgg actttctcaa gggggagaca ggcaagtacc tgcggctgcc  
 10 tcagctgggtg  
 1141 gacatggctg ctcatatgc ctcaggcatg gcgtacgtgg agcggatgaa  
 ctacgtccac  
 1201 cgggaccttc gtgcagccaa catcctgggtg ggagagaacc tgggtgtgcaa  
 agtggccgac  
 15 1261 tttgggctgg ctcggtcat tgaagacaat gactacacgg cgcggcaagg  
 tgccaaattc  
 1321 cccatcaagt ggacggctcc agaagctgcc ctctatggcc gcttcaccat  
 caagtcggac  
 1381 gtgtgggtcct tcgggatacct gctgactgag ctaccacaa agggacgggt  
 20 gccctaccct  
 1441 gggatgggtga accgcgaggt gctggaccag gtggagcggg gctaccggat  
 gccctgcccg  
 1501 ccggagtgtc ccgagtcct gcacgacctc atgtgccagt gctggcggaa  
 ggagcctgag  
 25 1561 gagcggccca ccttcagata cctgcaggcc ttctggagg actacttcac  
 gtccaccgag  
 1621 cccagtagc agcccgggga gaacctctag gcacaggcgg gccagaccg  
 gcttctcggc  
 1681 ttggatcctg ggctgggtgg cccctgtctc ggggcttgcc ccaactctgcc  
 30 tgctgtgtg  
 1741 tggctctctc tctgtggggc tgaattgcca ggggcgaggc cttctctct  
 tgggtggcatg  
 1801 gaaggggctt ctggacctag ggtggcctga gagggcgggtg ggtatgcgag  
 accagcacgg  
 35 1861 tgactctgtc cagctcccgc tgtggccgca cgcctctccc tgactcctc  
 cctggagctc  
 1921 tgtgggtctc tggaagagga accaggagaa gggctggggc cggggctgag  
 ggtgcccttt

1981 tccagcctca gcctactcgg ctcaactgaac tccttcccc cttctgtgcc  
 acccccggtc  
 2041 tatgtcgaga gctggccaaa gagcctttcc aaagaggagc gatgggcccc  
 tggccccgcc  
 5 2101 tgctgccac cctgccccct gccatccatt ctggaaacac ctgtaggcag  
 aggctgccga  
 2161 gacagaccct ctgcgcgtgc ttccaggctg ggcagcacia ggccttgctt  
 ggcctgatga  
 2221 tgggtgggtgg gtgggatgag taccacctca aacctgccc tccttagacc  
 10 tgagggaacc  
 2281 ttcgagatca tcacttcctt gcccccatth caccatggg gagacagttg  
 agagcgggga  
 2341 tgtgacatgc ccaaggccac ggagcagttc agagtggagg cgggcttggg  
 acccggtgct  
 15 2401 ccctctgtca tcctcaggaa ccaacaattc gtcggaggca tcatggaaag  
 actgggacag  
 2461 cccaggaaac aaggggtctg aggatgcatt cgagatggca gattccact  
 gccgctgccc  
 2521 gctcagccca gctgttggga acagcatgga ggcagatgtg gggctgagct  
 20 ggggaatcag  
 2581 ggtaaaaggt gcaggtgtgg agagagaggg ttcaatcggc ttgtgggtga  
 tgtttgacct  
 2641 tcagagccag ccggctatga aaggagcga gccccctggc tctggaggca  
 atcaagcaga  
 25 2701 catagaagag ccaagagtc aggaggccct ggtcctggcc tccttccccg  
 tactttgtcc  
 2761 cgtggcattt caattcctgg ccctgttctc ctccccaagt cggcaccctt  
 taactcatga  
 2821 ggagggaaaa gagtgcctaa gcggggggtga aagaggacgt gttaccact  
 30 gccatgcacc  
 2881 aggactggct gtgtaacctt ggggtggccc tgctgtctct ctgggctgca  
 gagtctgccc  
 2941 cacatgtggc catggcctct gcaactgtc agctctggc caggccctgt  
 ggcaggacac  
 35 3001 acatggtgag cctagccctg ggacatcagg agactgggct ctggctctgt  
 tcggcctttg  
 3061 ggtgtgtggt ggattctccc tgggcctcag tgtgcccac tgtaaagggg  
 cagctgacag

3121 tttgtggcat cttgccaagg gtccctgtgt- gtgtgtatgt gtgtgcatgt  
 gtgcgtgtct  
 3181 ccatgtgcgt ccatatttaa catgtaaaaa tgtccccccc gtcctgcccc  
 ccaaacatgt  
 5 3241 tgtacatttc accatggccc cctcatcata gcaataacat tcccactgcc  
 aggggttctt  
 3301 gagccagcca ggccctgcc gtggggaagg aggccaagca gtgcctgcct  
 atgaaatttc  
 3361 aacttttctt ttcatacgtc tttattaccc aagtcttctc ccgtccattc  
 10 cagtcaaata  
 3421 tgggctcact caccacagcg agctctcaaa tccctctcca actgcctaag  
 gccctttgtg  
 3481 taagggtgtc taatactgtc cttttttttt ttttaacagt gttttgtaga  
 tttcagatga  
 15 3541 ctatgcagag gcctggggga cccctggctc tgggccgggc ctggggctcc  
 gaaattccaa  
 3601 ggcccagact tgcggggggg ggggggggtat ccagaattgg ttgtaaatac  
 tttgcattat  
 3661 gtctgattaa acacaaacag acctcagaaa aaaaaaaaaa aaaaaaaaaa a  
 20

Table 5:

25 Mouse Src cDNA sequence (SEQ ID NO. 5)

1 atgggcagca acaagagcaa gcccaaggac gccagccagc ggccgagcag  
 cctggagccc  
 61 tcggaaaacg tgcacggggc agggggcgcc ttccggcct cacagacacc  
 30 gagcaagccc  
 121 gcctccgccc acggccaccg cgggccagc gccgccttcg tgccgcccgc  
 ggccgagccc  
 181 aagctcttcg gaggtttcaa ctctcggac accgtcacct ccccgagag  
 ggcgggogct  
 35 241 ctggcagggtg gggtagaccac ctttgtggcc ctctatgact atgagtcag  
 gacagagact  
 301 gacctgtcct tcaagaaagg ggagcggctg cagattgtca ataacacgag  
 gaagggtgat



361 gtcagagagg gagactggtg gctggcacac togetgagca cgggacagac  
 cggttacatc  
 421 cccagcaact atgtggcgcc ctccgactcc atccaggctg aggagtggta  
 ctttggcaag  
 5 481 atcactagac gggaaatcaga gcggtgctg ctcaacgccg agaaccgag  
 agggaccttc  
 541 ctctgaggg agagtgagac cacaaaaggt gcctactgcc tctctgtatc  
 cgacttcgac  
 601 aatgccaagg gcctaaatgt gaaacactac aagatccgca agctggacag  
 10 cggcggtttc  
 661 tacatcacct cccgcacca gttcaacagc ctgcagcagc tctggtgcta  
 ctactccaaa  
 721 catgtgatg gcctgtgtca ccgcctcact accgtatgtc ccacatccaa  
 gcctcagacc  
 15 781 cagggattgg ccaaggatgc gtgggagatc ccccgaggat ccttgcggct  
 ggaggtcaag  
 841 ctgggccagg gttgcttcgg agaggtgtgg atggggacct ggaacggcac  
 cagagggtt  
 901 gccatcaaaa ctctgaagcc aggcaccatg tcccagagg ccttctgca  
 20 ggaggcccaa  
 961 gtcataaga aactgaggca cgagaaactg gtgcagctgt atgtgtggt  
 gtcggaagaa  
 1021 ccatttaca ttgtgacaga gtacatgaac aaggggagtc tgctggactt  
 tctcaagggg  
 25 1081 gaaacgggca aatatttgcg gctacccag ctggtggaca tgtctgtca  
 gatcgcttca  
 1141 ggcattgctt atgtggagcg gatgaactat gtgcaccggg accttcgagc  
 cgccaatatc  
 1201 ctatgagggg agaacctggt gtgcaaagt ggcgactttg ggttgccccg  
 30 gctcatagaa  
 1261 gacaacgaat acacagccc gcaagggtgc aaattcccca tcaagtggac  
 cgccctgaa  
 1321 gctgtctgt acggcaggtt caccatcaag tcggatgtgt ggtcctttgg  
 gattctgtg  
 35 1381 accgagctca ccactaaggg aagagtgcc tatcctggga tggatgaaccg  
 tgaggttctg  
 1441 gaccaggtgg agcggggcta ccgatgcct tgcaccccg agtgcaccca  
 gtccttgcac

1501 gaccttatgt gccagtgtg gcggaaggag cccgaggagc ggcccacctt  
cgagtacctg

1561 caggccttcc tggaagacta ctttacgtcc actgagccac agtaccagcc  
cggggagaaac

5 1621 ctatag

## Claims

1. Method for identifying a therapeutic compound for the treatment of Alzheimer's disease comprising the steps
  - a) providing a Src protein and
  - b) determining the inhibitory effect of a compound on the Src activity.
2. Method of identifying a therapeutic compound for the treatment of Alzheimer's disease comprising the steps
  - a) providing a sequence which regulates Src expression and
  - b) determining if a compound inhibits the expression of Src protein.
3. Method according to claim 1 or 2, wherein Src is human Src or mouse Src.
4. Method according to one of the forgoing claims, wherein Src has the sequence SEQ ID NO. 1, SEQ ID NO. 2 or SEQ ID NO. 3.
5. Method according to claim 1, wherein Src is expressed in a mammalian cell.
6. Method according to claim 2, wherein the regulating sequence is the Src promotor.
7. Method of preparing a pharmaceutical for the treatment of Alzheimer's disease comprising the steps
  - a) identification of a therapeutic compound by using one of the methods claimed in claims 1 to 6;

- b) optimisation of the identified compound; and
- c) formulation of the optimised compound.

5 8. Use of a compound identified by a method according to claims 1 to 6 for the preparation of a pharmaceutical for treating Alzheimer's disease.

9. PP2 for use as pharmaceutical.

10 10. Use of PP2 for the preparation of a pharmaceutical for the treatment of Alzheimer's disease.

15

FRAV2002/0030

5

PATENT APPLICATION

10

INHIBITORS OF SRC KINASE FOR USE IN ALZHEIMER'S DISEASE

15

AVENTIS PHARMA S.A.

20

ABSTRACT

The present invention relates to the identification of inhibitors of Src kinase, assays and methods useful for such identification and the use of Src inhibitors for the preparation of pharmaceuticals for the treatment of Alzheimer's disease.

**THIS PAGE BLANK (USPTO)**

## 1. Figure 1




Figure 2







Figure 3

Figure 3 A



5 Figure 3 B





Figure 4



\*  $P < 0.01$

Figure 5





Figure 6



## Organization Applicant

-----  
 Street : 20, Avenue Raymond Aron  
 City : Antony  
 State : 92160  
 Country :  
 PostalCode :  
 PhoneNumber : 0033 1 55 71 4457  
 FaxNumber : 0033 1 5571 6172  
 EmailAddress : Claudia.Meinken@Aventis.com  
 <110> OrganizationName : Aventis Pharma S. A.

## Application Project

-----  
 <120> Title : Inhibitors of Src kinase for use in Alzheimer's disease  
 <130> AppFileReference : FRAV2002/0030  
 <140> CurrentAppNumber :  
 <141> CurrentFilingDate : \_\_\_\_-\_\_\_\_-\_\_\_\_

## Sequence

-----  
 <213> OrganismName : Homo sapiens  
 <400> PreSequenceString :  
 MGSNKS KPKD ASQRRRSLEP AENVHGAGGG AFPASQTPSK PASADGHRGP SAAFAPAAAE 60  
 PKLFGGFNSS DVTSPQRAG PLAGGVTTFFV ALYDYESRTE TDLSFKKGER LQIVNNTRKV 120  
 DVREGDWWLA HSLSTGQTGY IPSNYVAPSD SIQAEWYFG KITRRESERL LLNAENPRGT 180  
 FLVRESETTK GAYCLSVSDF DNAKGLNVKH YKIRKLDSGG FYITSRTQFN SLQQLVAYYS 240  
 KHADGLCHRL TTVCPSTSKPQ TQGLAKDAWE IPRESLRLEV KLGQGC FGEV WMGTWNGTTR 300  
 VAIKTLKPGT MSPEAFLQEA QVMKKLRHEK LVQLYAVVSE EPIYIVTEYM SKGSLLDFLK 360  
 GETGKYLRLP QLVDMAAQIA SGMAYVERMN YVHRDLRAAN ILVGENLVCK VADFGRLRLI 420  
 EDNEYTARQG AKFPIKWTAP EAALYGRFTI KSDVWSFGIL LTELTTKGRV PYPGMVNREV 480  
 LDQVERGYRM PCPPECPESL HDLMCQCWRK EPEERPTFEY LQAFLEDYFT STEPQYQPGE 540  
 NL 542  
 <212> Type : PRT  
 <211> Length : 542  
 SequenceName : human src isoform 1  
 SequenceDescription :

## Feature

-----  
 Sequence: human src isoform 1:  
 <221> FeatureKey : PEPTIDE  
 <222> LocationFrom : 1  
 <222> LocationTo : 542  
 Other Information :  
 CDSJoin : No

## Sequence

-----  
 <213> OrganismName : Homo sapiens  
 <400> PreSequenceString :  
 MGSNKS KPKD ASQRRRSLEP AENVHGAGGG AFPASQTPSK PASADGHRGP SAAFAPAAAE 60  
 PKLFGGFNSS DVTSPQRAG PLAGGVTTFFV ALYDYESRTE TDLSFKKGER LQIVNNTEGD 120  
 WWLAHSLSTG QTGYIPSNYV APSDSIQAE WYFGKITRRE SERLLLNAEN PRGTFLVRES 180  
 ETTKGAYCLS VSDFDNAKGL NVKHYKIRKL DSGGFYITSR TQFNSLQQLV AYYSKHADGL 240

```

src.WorkFile.txt
CHRLTTVCPT SKPQTQGLAK DAWEIPRESL RLEVKLGGQC FGEVWMGTWN GTTRVAIKTL 300
KPGTMSPEAF LQEAQVMKKL RHEKLVQLYA VVSEEPYIV TEYMSKGSLL DFLKGETGKY 360
LRLPQLVDMA AQIASGMAYV ERMNYVHRDL RAANILVGEN LVCKVADFGL ARLIEDNEYT 420
ARQGAKFPIK WTAPEAAALYG RFTIKSDVWS FGILLTELTT KGRVPYPGMV NREVLDQVER 480
GYRMPCPPEC PESLHDLMCQ CWRKEPEERP TFEYLQAFLE DYFTSTEPQY QPGENL 536

```

```

<212> Type : PRT
<211> Length : 536
      SequenceName : human src isoform 2
      SequenceDescription :

```

#### Feature

-----  
Sequence: human src isoform 2:

<221> FeatureKey : PEPTIDE

<222> LocationFrom : 1

<222> LocationTo : 536

Other Information :

CDSJoin : No

#### Sequence

-----  
<213> OrganismName : Murinae gen. sp.

<400> PreSequenceString :

```

MGSNKS KPKD ASQRRRSLEP SENVHGAGGA FPASQTPSKP ASADGHRGPS AAFVPPAAEP 60
KLFGGFNSSD TVTSPQRAGA LAGGVTTFVA LYDYESRTET DLSFKKGERL QIVNNTRKVD 120
VREGDWL LAH SLSTGQTGYI PSNYVAPSDS IQAEWYFGK ITRRESERLL LNAENPRGTF 180
LVRESETTKG AYCLSVSDFD NAKGLNVKHY KIRKLDSGGF YITSRTQFNS LQQLVAYYSK 240
HADGLCHRLT TVCPTSKPQT QGLAKDAWEI PRESLRLEV KLGQCFGEVW MGTWNGTTRV 300
AIKTLKPGTM SPEAFLQEAQ VMKKLRHEKL VQLYAVVSEE PIYIVTEYMN KGSLLDFLKG 360
ETGKYLRLPQ LVDMSAQIAS GMAYVERMNY VHRDLRAANI LVGENLVCKV ADFGLARLIE 420
DNEYTARQGA KFPIKWTAPE AALYGRFTIK SDVWSFGILL TELTTKGRVP YPGMVNREVL 480
DQVERGYRMP CPPECPESLH DLMCQCWRKE PEERPTFEYL QAFLEDYFTS TEPQYQPGEN 540
L 541

```

```

<212> Type : PRT
<211> Length : 541
      SequenceName : mouse scr
      SequenceDescription :

```

#### Feature

-----  
Sequence: mouse scr:

<221> FeatureKey : PEPTIDE

<222> LocationFrom : 1

<222> LocationTo : 541

Other Information :

CDSJoin : No

#### Sequence

-----  
<213> OrganismName : Homo sapiens

<400> PreSequenceString :

```

catcgaggtt ttgagaggct aactctccca aaaaggacca tgggtagcaa caagagcaag 60

```

cccaaggatg ccagccagcg gcgccgcagc ctggagcccg ccgagaacgt gcacggcgct	120
ggcgggggcg ctttccccgc ctgcagacc ccagcaagc cagcctcggc cgacggccac	180
cgcgggccca gcgcggcctt cgcccccgcg gccgccgagc ccaagctgtt cggaggcttc	240
aactcctcgg acaccgtcac ctccccgcag agggcgggcc cgctggccgg tggagtgacc	300
acctttgtgg ccctctatga ctatgagtct aggacggaga cagacctgtc cttcaagaaa	360
ggcgagcggc tccagattgt caacaacaca gaggagact ggtggctggc ccactcgctc	420
agcacaggac agacaggcta catccccagc aactacgtgg cgccctccga ctccatccag	480
gctgaggagt ggtatttttg caagatcacc agacgggagt cagagcgggt actgctcaat	540
gcagagaacc cgagagggac cttcctcgtg cgagaaagt agaccacgaa aggtgcctac	600
tgcctctcag tgtctgactt cgacaacgcc aagggcctca acgtgaagca ctacaagatc	660
cgcaagctgg acagcggcgg cttctacatc acctcccgca ccagttcaa cagcctgcag	720
cagctggtgg cctactactc caaacacgcc gatggcctgt gccaccgcct caccaccgtg	780
tgccccacgt ccaagccgca gactcagggc ctggccaagg atgcctggga gatccctcgg	840
gagtcgctgc ggctggaggt caagctgggc cagggtgct ttggcgaggt gtggatgggg	900
acctggaacg gtaccaccag ggtggccatc aaaacctga agcctggcac gatgtctcca	960
gaggccttcc tgcaggaggc ccaggatcat aagaagctga ggcatgagaa gctggtgcag	1020
ttgtatgctg tggtttcaga ggagcccatt tacatcgtca cggagtacat gagcaagggg	1080
agtttgctgg actttctcaa gggggagaca ggcaagtacc tgcggtgcc tcagctggtg	1140
gacatggctg ctcatatcgc ctcatgcatg gcgtacgtgg agcggatgaa ctacgtccac	1200
cgggaccttc gtgcagccaa catcctgggt ggagagaacc tgggtgtgcaa agtggccgac	1260
tttgggctgg ctcggtcat tgaagacaat gagtacacgg cgcggaagg tgccaaattc	1320
cccatcaagt ggacggctcc agaagctgcc ctctatggcc gcttcacat caagtggac	1380
gtgtggtcct tcgggacatc gctgactgag ctaccacaa agggacgggt gccctaccct	1440
gggatggtga accgcgaggt gctggaccag gtggagcggg gctaccggat gccctgcccg	1500
ccggagtgtc ccgagtcct gcacgacctc atgtgccagt gctggcggaa ggagcctgag	1560
gagcgccca cttcgagta cctgcaggcc ttcctggagg actacttcac gtccaccgag	1620
ccccagtacc agcccgggga gaacctctag gcacaggcgg gccagaccg gcttctcggc	1680
ttggatcctg ggctgggtgg cccctgtctc ggggcttgcc ccactctgcc tgcctgctgt	1740
tggtcctctc tctgtggggc tgaattgcc ggggcgaggc cttcctctt tgggtggcatg	1800
gaaggggctt ctggacctag ggtggcctga gaggcggtg ggtatgcgag accagcacgg	1860
tgactctgtc cagctcccgc tgtggccgca cgctctccc tgcactccct cctggagctc	1920
tgtgggtctc tggaagagga accaggagaa gggctggggc cgggctgag ggtgcccttt	1980
tccagcctca gcctactccg ctactgaac tccttccca cttctgtgcc acccccggtc	2040
tatgtcgaga gctggccaaa gagcctttcc aaagaggagc gatgggcccc tggccccgcc	2100

```

src.WorkFile.txt
tgcttgccac cctgccccctt gccatccatt ctggaaacac ctgtaggcag aggctgccga 2160
gacagaccct ctgccgctgc ttccaggctg ggcagacaaa ggccttgccct ggcctgatga 2220
tggtgggtgg gtgggatgag taccacctca aacctgccc tccttagacc tgagggaccc 2280
ttcgagatca tactttcctt gcccccatth caccatggg gagacagttg agagcgggga 2340
tgtgacatgc ccaaggccac ggagcagttc agagtggagg cgggcttgga acccggtgct 2400
ccctctgtca tcctcaggaa ccaacaattc gtcggaggca tcatggaaag actgggacag 2460
cccaggaaac aaggggtctg aggatgcatt cgagatggca gattcccact gccgctgccc 2520
gctcagccca gctgttgga acagcatgga ggcagatgtg gggctgagct ggggaatcag 2580
ggtaaaaggt gcaggtgtgg agagagaggc ttcaatcggc ttgtgggtga tgtttgacct 2640
tcagagccag ccggctatga aaggagcga gcccctcggc tctggaggca atcaagcaga 2700
catagaagag ccaagagtcc aggaggccct ggtcctggcc tccttccccg tactttgtcc 2760
cgtggcattt caattcctgg ccctgttctc ctcccaagt cggcaccctt taactcatga 2820
ggagggaaaa gagtgcctaa gcgggggtga aagaggacgt gttaccact gccatgcacc 2880
aggactggct gtgtaacctt gggtgccccc tgctgtctct ctgggctgca gagtctgccc 2940
cacatgtggc catggcctct gcaactgctc agctctggc caggccctgt ggcaggacac 3000
acatggtag cctagccctg ggacatcagg agactgggct ctggctctgt tcggcctttg 3060
gggtgtggtt ggatttctcc tgggcctcag tgtgccatc tgtaaagggg cagctgacag 3120
tttgtggcat cttgccaaag gtccctgtgt gtgtgtatgt gtgtgcatgt gtgctgtct 3180
ccatgtgcgt ccatatttaa catgtaaaaa tgtcccccc gctccgtccc ccaaacatgt 3240
tgtacatttc accatggccc cctcatcata gcaataacat tcccactgcc aggggttctt 3300
gagccagcca ggcctgcca gtggggaagg aggccaaagca gtgcctgcct atgaaatttc 3360
aacttttctt ttcatactgc tttattacc aagtcttctc ccgtccattc cagtcaaadc 3420
tggtgtcact caccacagcg agctctcaaa tccctctcca actgcctaag gccctttgtg 3480
taaggtgtct taatactgtc cttttttttt ttttaacagt gttttgtaga tttcagatga 3540
ctatgcagag gcctggggga cccctggctc tgggccgggc ctggggctcc gaaattccaa 3600
ggcccagact tgcggggggg ggggggggtat ccagaattgg ttgtaaatac tttgcatatt 3660
gtctgattaa acacaaacag acctcagaaa aaaaaaaaaa aaaaaaaaaa a 3711

```

```

<212> Type : DNA
<211> Length : 3711
      SequenceName : human src
      SequenceDescription :

```

#### Feature

```

-----
Sequence: human src:
<221> FeatureKey : CDS
<222> LocationFrom : 1
<222> LocationTo : 3711
      Other Information :
      CDSJoin : No

```



## Custom Codon

Sequence Name : human src

## Sequence

&lt;213&gt; OrganismName : Murinae gen. sp.

&lt;400&gt; PreSequenceString :

```

atgggcagca acaagagcaa gcccaaggac gccagccagc ggcgccgcag cctggagccc      60
tcggaaaacg tgcacggggc agggggcgcc ttcccggcct cacagacacc gagcaagccc      120
gcctccgccg acggccaccg cgggcccagc gccgccttcg tgccgcccgc ggccgagccc      180
aagctcttcg gaggcttcaa ctctcggac accgtcacct ccccgagag ggcgggcgct      240
ctggcaggtg gggtgaccac ctttgtggcc ctctatgact atgagtcacg gacagagact      300
gacctgtcct tcaagaaagg ggagcggctg cagattgtca ataacacgag gaagggtgat      360
gtcagagagg gagactgggt gctggcacac tcgctgagca cgggacagac cggttacatc      420
cccagcaact atgtggcgcc ctccgactcc atccaggctg aggagtggta ctttggcaag      480
atcactagac gggaaatcaga gcggctgctg ctcaacgccg agaaccgag agggaccttc      540
ctcgtgaggg agagtgagac caaaaagggt gcctactgcc tctctgtatc cgacttcgac      600
aatgccaagg gcctaaatgt gaaacactac aagatccgca agctggacag cggcgggtttc      660
tacatcacct cccgcaccca gttcaacagc ctgcagcagc tcgtggctta ctactccaaa      720
catgctgatg gcctgtgtca ccgcctcact accgtatgtc ccacatccaa gcctcagacc      780
cagggatttg ccaaggatgc gtgggagatc ccccgaggat ccctgcggct ggagggtcaag      840
ctgggccagg gttgcttcgg agagggtgtg atggggacct ggaacggcac cagagggtt      900
gccatcaaaa ctctgaagcc aggcaccatg tcccagagg ccttcctgca ggaggcccaa      960
gtcatgaaga aactgaggca cgagaaactg gtgcagctgt atgctgtggt gtcggaagaa     1020
cccatttaca ttgtgacaga gtacatgaac aaggggagtc tgctggactt tctcaagggg     1080
gaaacgggca aatattttcg gtagccccag ctggtggaca tgtctgctca gatcgttca     1140
ggcatggcct atgtggagcg gatgaactat gtgcaccggg accttcgagc cgccaatatc     1200
ctagtagggg agaacctggt gtgcaaagtg gccgactttg ggttggcccc gctcatagaa     1260
gacaacgaat acacagcccc gcaagggtgcc aaattcccca tcaagtggac cgcccctgaa     1320
gctgctctgt acggcagggt caccatcaag tcggatgtgt ggtcctttgg gattctgctg     1380
accgagctca ccactaaggg aagagtgcc tatcctggga tggatgaacc tgagggtctg     1440
gaccaggtgg agcggggcta ccggatgcct tgtcccccg agtgccccga gtccctgcat     1500
gaccttatgt gccagtgtg gcggaaggag cccgaggagc ggcccacctt cgagtacctg     1560
caggccttcc tggaagacta ctttacgtcc actgagccac agtaccagcc cggggagaa      1620
ctatag

```

&lt;212&gt; Type : DNA

&lt;211&gt; Length : 1626

SequenceName : Mouse src

SequenceDescription :

src.WorkFile.txt

Feature

-----

Sequence: Mouse src:  
<221> FeatureKey : CDS  
<222> LocationFrom : 1  
<222> LocationTo : 1626  
Other Information :  
CDSJoin : No

Custom Codon

-----

Sequence Name : Mouse src